

BOUNDS FOR THE ENTROPY OF GRADED ALGEBRAS

JAN SNELLMAN

ABSTRACT. Newman, Schneider and Shalev defined the *entropy* of a graded associative algebra A as $H(A) = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}$, where a_n is the vector space dimension of the n 'th homogeneous component. When A is the homogeneous quotient of a finitely generated free associative algebra, they showed that $H(A) \leq \sqrt{a_2}$. Using some results of Friedland on the maximal spectral radius of $(0, 1)$ -matrices with a prescribed number of ones, we improve on this bound.

Let $A = \bigoplus_{n=0}^{\infty} A_n$ be a graded, associative algebra over a field F . Assume furthermore that A is infinite-dimensional as a vector space over F , but that each graded component A_n is finite-dimensional, so that $a_n = \dim A_n < \infty$. Newman, Schneider and Shalev [3] defined the *entropy* of A by

$$H(A) = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}. \quad (1)$$

Clearly, if a_n has polynomial growth, then $H(A) = 1$, and if a_n has exponential growth, with $a_n \sim cd^n$, then $H(A) = d$. Denote the Hilbert series of A by

$$A(t) = \sum_{n=0}^{\infty} a_n t^n. \quad (2)$$

Then $H(A) = 1/R$, where R is the radius of convergence of $A(t)$. If the Hilbert series is rational with $A(t) = P(t)/Q(t)$, then $|\lambda_{\min}| \leq H(A) \leq |\lambda_{\max}|$, where $\lambda_{\min}$ and $\lambda_{\max}$ are roots of $Q(t)$ with minimal and maximal modulus.

We henceforth assume that A is the quotient $A = F[X^*]/I$, where $F[X^*]$ is the free associative algebra on a finite alphabet X , and I is a two-sided homogeneous ideal. Then A is connected, i.e. $A_0 = F$, and generated by A_1 . Newman, Schneider and Shalev observed that the series $(a_n)_{n=0}^{\infty}$ is submultiplicative, i.e. $a_n a_m \leq a_{n+m}$, and that this implies that

- $\lim_{n \rightarrow \infty} \sqrt[n]{a_n}$ exists and is bounded by $\sqrt[n]{a_n}$ for each $n \geq 1$, and consequently that
- $H(A) \leq \sqrt[n]{a_n}$ for all $n \geq 1$.

In particular,

$$H(A) \leq \sqrt{a_2}, \quad (3)$$

and this bound is obtained when $a_2 = m^2$ and A is the free associative algebra on m letters.

Using a graph-theoretical result of Friedland [2] and Brualdi & Hoffman [1] we can improve on this bound:

1991 *Mathematics Subject Classification*. 16W50, 05C50; 16P90, 05C20, 05C38.

Key words and phrases. Entropy of algebras, spectral radius, digraphs, 0 – 1 matrices, Perron-Frobenius theorem.

Theorem 0.1. *Let $a_2 = k$, with*

$$k = m^2 + \ell, \quad 1 \leq \ell \leq 2m.$$

Then

$$H(A) \leq \frac{m + \sqrt{m^2 + 2\ell}}{2} =: f(k) \quad (4)$$

If $\ell = 1$ then

$$H(A) \leq m, \quad (5)$$

if $\ell = 2m - 3$ then

$$H(A) \leq \frac{m - 1 + \sqrt{m^2 + 6m - 7}}{2}. \quad (6)$$

Let $\rho_{m,p,q}$ denote the largest positive root of

$$t^3 - mt^2 - pt + mp - pq \quad (7)$$

Then, for any ℓ , there is an $M(\ell)$ such that for $m \geq M(\ell)$ we have that

$$\dim A_2 = m^2 + \ell \implies H(A) \leq \rho_{m, \lfloor \ell/2 \rfloor, \lceil \ell/2 \rceil} \quad (8)$$

Proof. Recall that we've assumed that $A = F[X^*]/I$. Denote by d the number of letters in X , so that $X = \{x_1, x_2, \dots, x_d\}$. Let $\succ$ be a term-order on the free monoid X^* , and let $B = F[X^*]/\text{in}(I)$, where $\text{in}(I)$ is the *initial ideal* of I w.r.t $\succ$. Since $A(t) = B(t)$ we have that $H(A) = H(B)$. Now, if we let J denote the monomial ideal generated by the quadratic part of $\text{in}(I)$, then $C = F[X^*]/J$ is a quadratic monomial algebra. Clearly $a_2 = b_2 = c_2$, and $A(t) = B(t) \ll C(t)$, i.e. $C(t) = \sum_{n=0}^{\infty} c_n t^n$ and $c_n \geq a_n = b_n$ for all n . It follows that $H(A) = H(B) \leq H(C)$.

Let G be the directed graph with vertex set $\{1, 2, \dots, d\}$, and with an (directed) edge from i to j iff $x_i x_j \notin J$. Thus, we allow loops, but no multiple edges. Clearly, for $n \geq 2$ there is a bijection between on the one hand the set of those monomials in X^* that are of length n , and which do not belong to J , and on the other hand the set of directed walks in G of length n . Thus, if g_n denotes the number of directed walks in G of length n , then $g_n = c_n$.

Let M denote the adjacency matrix of G , i.e. that $d \times d$ matrix (m_{ij}) with $m_{ij} = 1$ iff there is an edge from i to j . Let $\mathbf{e} = (1, 1, \dots, 1)$ denote the (column) vector consisting of d ones. Then $g_n = \mathbf{e}^t M^n \mathbf{e}$. If

$$\phi(\lambda) = \lambda^d - h_1 \lambda^{d-1} - h_2 \lambda^{d-2} - \dots - h_{d-1} \lambda - h_d$$

denotes the characteristic equation of M , we have, by the Cayley-Hamilton theorem, that $\phi(M) = 0$. Hence

$$M^d = h_1 M^{d-1} + h_2 M^{d-2} + \dots + h_{d-1} M + h_d I,$$

which by multiplication with M^s yields

$$M^{s+d} = h_1 M^{s+d-1} + h_2 M^{s+d-2} + \dots + h_{d-1} M^{s+1} + h_d M^s,$$

hence

$$\begin{aligned}
g_{s+d} &= \mathbf{e}^t M^{s+d} \mathbf{e} \\
&= h_1 \mathbf{e}^t M^{s+d-1} \mathbf{e} + h_2 \mathbf{e}^t M^{s+d-2} \mathbf{e} + \cdots + h_{d-1} \mathbf{e}^t M^{s+1} \mathbf{e} + h_d \mathbf{e}^t M^s \mathbf{e} \\
&= h_1 g_{s+d-1} + h_2 g_{s+d-2} + \cdots + h_{d-1} g_{s+1} + h_d g_s
\end{aligned} \tag{9}$$

Since the g_n 's and hence the c_n 's satisfy the linear recurrence (9), it follows (as mentioned in section 2 of [3]) that $H(C) \leq |\rho|$, where ρ is the maximal modulus of a root of the characteristic polynomial of M , i.e. the largest absolute value of an eigenvalue of M , i.e. the spectral radius of M (and by definition of G).

It follows that we can apply the bounds obtained by Friedland [2] and Brualdi & Hoffman [1] for the spectral radius of directed graphs with k edges to bound the entropy of C , and hence that of A .

Let $G_{m,p,q}$ be the directed graph on $\{1, \dots, m+1\}$, where there is an edge from i to j if $i, j \leq m$ or if $i \leq p$ and $j = m+1$ or if $i = m+1$ and $j \leq q$. Thus, $G_{m,p,q}$ has adjacency matrix

$$\left[\begin{array}{cccccc|c} & & & & & & 1 \\ & & & & & & 1 \\ & & & & & & \vdots \\ & & & & & & 1 \\ & & & & & & 0 \\ & & & & & & \vdots \\ & & & & & & 0 \\ \hline 1 & 1 & \cdots & 1 & 0 & \cdots & 0 \end{array} \right]$$

with p ones on the last row, q ones on the last column, and with $\mathbb{I}_m$ an $m \times m$ matrix of ones. It corresponds to the quadratic monomial algebra

$$A_{m,p,q} = \langle x_1, \dots, x_m \mid x_{m+1}^2; x_m x_{m+1}, \dots, x_{p+1} x_{m+1}; x_{m+1} x_m, \dots, x_{m+1} x_{q+1} \rangle.$$

Let $\rho_{m,p,q}$ denote the spectral radius of $G_{m,p,q}$, hence $\rho_{m,p,q} = H(A_{m,p,q})$. Friedland [2] showed that $\rho_{m,p,q}$ is the largest positive root of (7) and that for $k = m^2 + 1$, the graph $G_{m,0,1}$ has maximal spectral radius $\rho_{m,0,1} = m$ of all directed graphs with k edges (the later result was also obtained by Brualdi and Hoffman [1]). This gives (5). Friedland also showed that for $k = m^2 + 2m - 3 = (m+1)^2 - 4$ edges, the maximal spectral radius is obtained *not* by $G_{m,m-2,m-1}$ but by the graph on $m+1$ vertices where there is an edge from i to j if either $i < m$ or $j < m$; i.e. by the digraph with adjacency matrix

$$\left[\begin{array}{cccc|cc} & & & & 1 & 1 \\ & & & & 1 & 1 \\ & & & & \vdots & \vdots \\ & & & & 1 & 1 \\ \hline 1 & 1 & \cdots & 1 & 0 & 0 \\ 1 & 1 & \cdots & 1 & 0 & 0 \end{array} \right]$$

Friedland showed that this graph has spectral radius

$$\frac{m-1+\sqrt{m^2+6m-7}}{2},$$


so we have (6).

Friedland furthermore showed that for a fixed ℓ it holds that for sufficiently large m , if $k = m^2 + \ell$ then $G_{m, \lfloor \ell/2 \rfloor, \lceil \ell/2 \rceil}$ has the largest spectral radius of a directed graph with k edges. This gives (8). $\square$

We have that for $k = m^2 + \ell$,

$$\begin{aligned} f(k) < \sqrt{k} &\iff f(k)^2 < k &\iff \left(\frac{m + \sqrt{m^2 + 2\ell}}{2} \right)^2 < m^2 + \ell \\ &\iff m^2 + \ell - m\sqrt{m^2 + 2\ell} > 0 &\iff \ell > 0, \end{aligned}$$

so the bound of (4) is indeed an improvement for $\ell > 0$. The number $\rho_{m, \lfloor \ell/2 \rfloor, \lceil \ell/2 \rceil}$ is smaller still, as is shown in the picture below, which plots $\sqrt{k} - f(k)$ (in a solid line) and $\sqrt{k} - \rho_{m, \lfloor \ell/2 \rfloor, \lceil \ell/2 \rceil}$ (in a dotted line) for $5 \leq k \leq 85$.



REFERENCES

- [1] R. A. Brualdi and A. J. Hoffman. On the Spectral Radius of (0,1)-Matrices. *Linear Algebra and its Applications*, 65:133–146, 1985.
- [2] Schmel Friedland. The Maximal Eigenvalue of 0-1 Matrices with Prescribed Number of Ones. *Linear Algebra and its Applications*, 69:33–69, 1985.
- [3] M. F. Newman, Csaba Schneider, and Aner Shalev. The Entropy of Graded Algebras. *Journal of Algebra*, 223:85–100, 2000.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-10691 STOCKHOLM, SWEDEN

E-mail address: jans@matematik.su.se